

New England Biolabs Certificate of Analysis

Product Name: Nt.CviPII
Catalog Number: R0626S
Concentration: 2,000 U/ml
Unit Definition: One unit is defined as the amount of enzyme required to digest 1 µg of pUC19 DNA in CutSmart™ Buffer incubated for 1 hour at 37°C in a total reaction volume of 50 µl.
Packaging Lot Number: 10230768
Expiration Date: 02/2025
Storage Temperature: -20°C
Storage Conditions: 100 mM NaCl, 20 mM Tris-HCl (pH 8.0), 50% Glycerol
Specification Version: PS-R0626S/L v2.0


Nt.CviPII Component List			
NEB Part Number	Component Description	Lot Number	Individual QC Result
R0626SVIAL	Nt.CviPII	10226123	Pass
B6004SVIAL	rCutSmart™ Buffer	10224839	Pass

Assay Name/Specification	Lot # 10230768
DNase Activity (Labeled Oligo, 3' extension) A 50 µl reaction in CutSmart® Buffer containing a 20 nM solution of a fluorescent labeled double-stranded oligonucleotide containing a 3' extension and a minimum of 10 units of Nt.CviPII incubated for 16 hours at 37°C yields <5% degradation as determined by capillary electrophoresis.	Pass
DNase Activity (Labeled Oligo, 5' extension) A 50 µl reaction in CutSmart® Buffer containing a 20 nM solution of a fluorescent labeled double-stranded oligonucleotide containing a 5' extension and a minimum of 10 units of Nt.CviPII incubated for 16 hours at 37°C yields <5% degradation as determined by capillary electrophoresis.	Pass
Double Stranded DNase Activity (Labeled Oligo) A 50 µl reaction in CutSmart® Buffer containing a 20 nM solution of a fluorescent labeled double-stranded oligonucleotide containing a blunt end and a minimum of 10 units of Nt.CviPII incubated for 16 hours at 37°C yields <5% degradation as determined by capillary electrophoresis.	Pass
Single Stranded DNase Activity (FAM-Labeled Oligo)	Pass


Assay Name/Specification	Lot # 10230768
<p>A 50 µl reaction in CutSmart® Buffer containing a 20 nM solution of a fluorescent internal labeled oligonucleotide and a minimum of 10 units of Nt.CviPII incubated for 16 hours at 37°C yields <5% degradation as determined by capillary electrophoresis.</p> <p>qPCR DNA Contamination (E. coli Genomic) A minimum of 2 units of Nt.CviPII is screened for the presence of E. coli genomic DNA using SYBR® Green qPCR with primers specific for the E. coli 16S rRNA locus. Results are quantified using a standard curve generated from purified E. coli genomic DNA. The measured level of E. coli genomic DNA contamination is ≤ 1 E. coli genome.</p>	<p>Pass</p>

This product has been tested and shown to be in compliance with all specifications.

One or more products referenced in this document may be covered by a 3rd-party trademark. Please visit www.neb.com/trademarks for additional information.



Ana Egana
Production Scientist
05 Feb 2024



Michael Tonello
Packaging Quality Control Inspector
05 Feb 2024