

New England Biolabs Certificate of Analysis

Product Name: Nt.BbvCI
Catalog Number: R0632S
Concentration: 10,000 U/ml
Unit Definition: One unit is defined as the amount of enzyme required to convert 1 µg of supercoiled pUB DNA to open circular form in 1 hour at 37°C in a total reaction volume of 50 µl.
Lot Number: 10013100
Expiration Date: 06/2020
Storage Temperature: -20°C
Storage Conditions: 50 mM KCl, 10 mM Tris-HCl (pH 7.4), 1 mM DTT, 0.1 mM EDTA, 50% Glycerol, 200 µg/ml BSA
Specification Version: PS-R0632S/L v1.0

| Nt.BbvCI Component List | | | |
|-------------------------|-----------------------|------------|----------------------|
| NEB Part Number | Component Description | Lot Number | Individual QC Result |
| R0632SVIAL | Nt.BbvCI | 10012693 | Pass |
| B7204SVIAL | CutSmart® Buffer | 10014372 | Pass |

| Assay Name/Specification | Lot # 10013100 |
|--|----------------|
| Endonuclease Activity (Nicking) A 50 µl reaction in CutSmart™ Buffer containing 1 µg of supercoiled pUC19 DNA and a minimum of 30 units of Nt.BbvCI incubated for 4 hours at 37°C results in <10% conversion to the nicked form as determined by agarose gel electrophoresis. | Pass |
| Exonuclease Activity (Radioactivity Release) A 50 µl reaction in CutSmart™ Buffer containing 1 µg of a mixture of single and double-stranded [³ H] E. coli DNA and a minimum of 30 units of Nt.BbvCI incubated for 4 hours at 37°C releases <0.1% of the total radioactivity. | Pass |
| Non-Specific DNase Activity (16 Hour) A 50 µl reaction in CutSmart™ Buffer containing 1 µg of Lambda DNA and a minimum of 30 Units of Nt.BbvCI incubated for 16 hours at 37°C results in a DNA pattern free of detectable nuclease degradation as determined by agarose gel electrophoresis. | Pass |

This product has been tested and shown to be in compliance with all specifications.



Jianying Luo
Production Scientist
18 Jun 2018



Michael Tonello
Packaging Quality Control Inspector
15 Aug 2018