

## New England Biolabs Certificate of Analysis

**Product Name:** *Pcil*  
**Catalog Number:** *R0655L*  
**Concentration:** *10,000 U/ml*  
**Unit Definition:** *One unit is defined as the amount of enzyme required to digest 1 µg of pXba DNA in NEBuffer r3.1 in 1 hour at 37°C in a total reaction volume of 50 µl.*  
**Packaging Lot Number:** *10161996*  
**Expiration Date:** *08/2024*  
**Storage Temperature:** *-20°C*  
**Storage Conditions:** *10 mM Tris-HCl, 300 mM NaCl, 1 mM DTT, 0.1 mM EDTA, 50% Glycerol, 500 µg/ml rAlbumin (pH 7.4 @ 25°C)*  
**Specification Version:** *PS-R0655S/L v2.0*

Pcil Component List			
NEB Part Number	Component Description	Lot Number	Individual QC Result
R0655LVIAL	Pcil	10161999	Pass
B7024AVIAL	Gel Loading Dye, Purple (6X)	10156427	Pass
B6003SVIAL	NEBuffer™ r3.1	10146827	Pass

Assay Name/Specification	Lot # 10161996
<b>Protein Purity Assay (SDS-PAGE)</b> Pcil is ≥ 95% pure as determined by SDS-PAGE analysis using Coomassie Blue detection.	Pass
<b>Non-Specific DNase Activity (16 Hour)</b> A 50 µl reaction in NEBuffer™ r3.1 containing 1 µg of pXba DNA and a minimum of 50 units of Pcil incubated for 16 hours at 37°C results in a DNA pattern free of detectable nuclease degradation as determined by agarose gel electrophoresis.	Pass
<b>qPCR DNA Contamination (E. coli Genomic)</b> A minimum of 10 units of Pcil is screened for the presence of E. coli genomic DNA using SYBR® Green qPCR with primers specific for the E. coli 16S rRNA locus. Results are quantified using a standard curve generated from purified E. coli genomic DNA. The measured level of E. coli genomic DNA contamination is ≤ 1 E. coli genome.	Pass
<b>Exonuclease Activity (Radioactivity Release)</b> A 50 µl reaction in NEBuffer™ r3.1 containing 1 µg of a mixture of single and	Pass

Assay Name/Specification	Lot # 10161996
<p>double-stranded [<sup>3</sup>H] E. coli DNA and a minimum of 100 units of PciI incubated for 4 hours at 37°C releases &lt;0.1% of the total radioactivity.</p>	
<p><b>Endonuclease Activity (Nicking)</b> A 50 µl reaction in NEBuffer™ r3.1 containing 1 µg of supercoiled PhiX174 DNA and a minimum of 10 units of PciI incubated for 4 hours at 37°C results in &lt;20% conversion to the nicked form as determined by agarose gel electrophoresis.</p>	<b>Pass</b>
<p><b>Ligation and Recutting (Terminal Integrity)</b> After a 10-fold over-digestion of pXba DNA with PciI, &gt;95% of the DNA fragments can be ligated with T4 DNA ligase in 16 hours at 16°C. Of these ligated fragments, &gt;95% can be recut with PciI.</p>	<b>Pass</b>

This product has been tested and shown to be in compliance with all specifications.

One or more products referenced in this document may be covered by a 3rd-party trademark. Please visit [www.neb.com/trademarks](http://www.neb.com/trademarks) for additional information.



Penghua Zhang  
Production Scientist  
12 Sep 2022



Michael Tonello  
Packaging Quality Control Inspector  
11 Oct 2022