

240 County Road Ipswich, MA 01938-2723 Tel 978-927-5054 Fax 978-921-1350 www.neb.com info@neb.com

New England Biolabs Certificate of Analysis

Product Name: Zral

Catalog Number: R0659S

Concentration: 10,000 U/ml

Unit Definition: One unit is defined as the amount of enzyme required to digest 1 µg

Lambda DNA in rCutSmart™ Buffer in 1 hour at 37°C in a total

reaction volume of 50 µl.

Packaging Lot Number: 10134074
Expiration Date: 01/2024
Storage Temperature: -20°C

Storage Conditions: 10 mM Tris-HCl, 300 mM NaCl, 1 mM DTT, 0.1 mM EDTA, 50% Glycerol,

500 $\mu g/ml$ rAlbumin (pH 7.4 @ 25°C)

Specification Version: PS-R0659S/L/V v2.0

Zral Component List				
NEB Part Number	Component Description	Lot Number	Individual QC Result	
R0659SVIAL	Zral	10133995	Pass	
B6004SVIAL	rCutSmart™ Buffer	10127378	Pass	

Assay Name/Specification	Lot # 10134074
Protein Purity Assay (SDS-PAGE)	Pass
Zral is ≥ 95% pure as determined by SDS-PAGE analysis using Coomassie Blue detection.	
Ligation and Recutting (Terminal Integrity)	Pass
After a 10-fold over-digestion of Lambda DNA with Zral, >95% of the DNA fragments	
can be ligated with T4 DNA ligase in 16 hours at 16°C. Of these ligated fragments, >95% can be recut with Zral.	
Non-Specific DNase Activity (16 Hour)	Pass
A 50 µl reaction in rCutSmart™ Buffer containing 1 µg of Lambda DNA and a minimum of	
10 units of Zral incubated for 16 hours at 37°C results in a DNA pattern free of detectable nuclease degradation as determined by agarose gel electrophoresis.	
detectable nuclease degradation as determined by againse gerelectrophoresis.	
qPCR DNA Contamination (E. coli Genomic)	Pass
A minimum of 10 units of Zral is screened for the presence of E. coli genomic DNA	
using SYBR® Green qPCR with primers specific for the E. coli 16S rRNA locus. Results	
are quantified using a standard curve generated from purified E. coli genomic DNA.	



R0659S / Lot: 10134074

Page 1 of 2

Assay Name/Specification	Lot # 10134074
The measured level of E. coli genomic DNA contamination is ≤ 1 E. coli genome.	
Endonuclease Activity (Nicking)	Pass
A 50 µl reaction in rCutSmart™ Buffer containing 1 µg of supercoiled M13mp18 DNA and a minimum of 10 units of Zral incubated for 4 hours at 37°C results in <20%	
conversion to the nicked form as determined by agarose gel electrophoresis.	
Exonuclease Activity (Radioactivity Release)	Pass
A 50 µl reaction in rCutSmart™ Buffer containing 1 µg of a mixture of single and	
double-stranded [³H] E. coli DNA and a minimum of 100 units of Zral incubated for 4 hours at 37°C releases <0.1% of the total radioactivity.	

This product has been tested and shown to be in compliance with all specifications.

One or more products referenced in this document may be covered by a 3rd-party trademark. Please visit www.neb.com/trademarks for additional information.

Penghaa Zhang Production Scientist

04 Feb 2022

Josh Hersey

Packaging Quality Control Inspector

04 Feb 2022



R0659S / Lot: 10134074

Page 2 of 2