

New England Biolabs Certificate of Analysis

Product Name: BspQI
Catalog Number: R0712S
Concentration: 10,000 U/ml
Unit Definition: One unit is defined as the amount of enzyme required to digest 1 µg of Lambda DNA in NEBuffer r3.1 in 1 hour at 50°C in a total reaction volume of 50 µl.
Packaging Lot Number: 10233277
Expiration Date: 02/2026
Storage Temperature: -20°C
Storage Conditions: 10 mM Tris-HCl, 100 mM KCl, 1 mM DTT, 0.1 mM EDTA, 50% Glycerol, 200 µg/ml rAlbumin (pH 7.4 @ 25°C)
Specification Version: PS-R0712S/L v3.0

BspQI Component List			
NEB Part Number	Component Description	Lot Number	Individual QC Result
R0712SVIAL	BspQI	10228171	Pass
B6003SVIAL	NEBuffer™ r3.1	10227733	Pass

Assay Name/Specification	Lot # 10233277
Endonuclease Activity (Nicking) A 50 µl reaction in NEBuffer™ r3.1 containing 1 µg of supercoiled LITMUS38i DNA and a minimum of 10 units of BspQI incubated for 4 hours at 50°C results in <20% conversion to the nicked form as determined by agarose gel electrophoresis.	Pass
Exonuclease Activity (Radioactivity Release) A 50 µl reaction in NEBuffer™ r3.1 containing 1 µg of a mixture of single and double-stranded [³ H] E. coli DNA and a minimum of 50 units of BspQI incubated for 4 hours at 50°C releases <0.1% of the total radioactivity.	Pass
Functional Testing (15 minute Digest) A 50 µl reaction in NEBuffer™ r3.1 containing 1 µg of Lambda DNA and 1 µl of BspQI incubated for 15 minutes at 50°C results in complete digestion as determined by agarose gel electrophoresis.	Pass
Ligation and Recutting (Terminal Integrity) After a 10-fold over-digestion of Lambda DNA with BspQI, >95% of the DNA fragments can be ligated with T4 DNA ligase in 16 hours at 16°C. Of these ligated fragments,	Pass

Assay Name/Specification	Lot # 10233277
<p>>95% can be recut with BspQI.</p> <p>Non-Specific DNase Activity (16 hour) A 50 µl reaction in NEBuffer™ r3.1 containing 1 µg of Lambda DNA and a minimum of 10 units of BspQI incubated for 16 hours at 50°C results in a DNA pattern free of detectable nuclease degradation as determined by agarose gel electrophoresis. NOTE: although no nuclease degradation is detected under these conditions, extended incubations and/or high concentrations of this enzyme may result in star activity. See the product FAQ for recommended reaction conditions for this enzyme.</p>	Pass
<p>Protein Purity Assay (SDS-PAGE) BspQI is ≥ 95% pure as determined by SDS-PAGE analysis using Coomassie Blue detection.</p>	Pass
<p>RNase Activity (Extended Digestion) A 10 µl reaction in NEBuffer 4 containing 40 ng of a 300 base single-stranded RNA and a minimum of 10 units of BspQI is incubated at 37°C. After incubation for 4 hours, >90% of the substrate RNA remains intact as determined by gel electrophoresis using fluorescent detection.</p>	Pass
<p>qPCR DNA Contamination (E. coli Genomic) A minimum of 10 units of BspQI is screened for the presence of E. coli genomic DNA using SYBR® Green qPCR with primers specific for the E. coli 16S rRNA locus. Results are quantified using a standard curve generated from purified E. coli genomic DNA. The measured level of E. coli genomic DNA contamination is ≤ 1 E. coli genome.</p>	Pass

This product has been tested and shown to be in compliance with all specifications.

One or more products referenced in this document may be covered by a 3rd-party trademark. Please visit www.neb.com/trademarks for additional information.



Ana Egana
Production Scientist
23 Feb 2024



Josh Hersey
Packaging Quality Control Inspector
23 Feb 2024