

New England Biolabs Certificate of Analysis

Product Name: WarmStart® Nt.BstNBI
Catalog Number: R0725S
Concentration: 10,000 U/ml
Unit Definition: One unit is defined as the amount of enzyme required to digest 1 µg T7 DNA in NEBuffer r3.1 in 1 hour at 55°C in a total reaction volume of 50 µl.
Packaging Lot Number: 10154563
Expiration Date: 06/2024
Storage Temperature: -20°C
Storage Conditions: 10 mM Tris-HCl, 50 mM KCl, 1 mM DTT, 0.1 mM EDTA, 50% Glycerol, 200 µg/ml rAlbumin (pH 7.4 @ 25°C)
Specification Version: PS-R0725S v1.0

WarmStart® Nt.BstNBI Component List			
NEB Part Number	Component Description	Lot Number	Individual QC Result
R0725SVIAL	WarmStart® Nt.BstNBI	10154579	Pass
B6003SVIAL	NEBuffer™ r3.1	10146824	Pass

Assay Name/Specification	Lot # 10154563
Exonuclease Activity (Radioactivity Release) A 50 µl reaction in NEBuffer™ r3.1 containing 1 µg of a mixture of single and double-stranded [³ H] E. coli DNA and a minimum of 50 units of WarmStart® Nt.BstNBI incubated for 4 hours at 55°C releases <0.1% of the total radioactivity.	Pass
Functional Testing (WarmStart Inhibition) A 50 µl reaction in NEBuffer™ r3.1 containing 1 µg of T7 DNA and a minimum of 10 units of WarmStart® Nt.BstNBI incubated for 1 hour at 25°C results in <5% digestion of the DNA as determined by agarose gel electrophoresis.	Pass
Protein Purity Assay (SDS-PAGE) Nt.BstNBI is ≥ 95% pure as determined by SDS-PAGE analysis using Coomassie Blue detection.	Pass
Non-Specific DNase Activity (16 hour) A 50 µl reaction in NEBuffer™ r3.1 containing 1 µg of T7 DNA and a minimum of 10 units of WarmStart® Nt.BstNBI incubated for 16 hours at 55°C results in a DNA pattern free of detectable nuclease degradation as determined by agarose gel	Pass

Assay Name/Specification	Lot # 10154563
<p>electrophoresis. NOTE: although no nuclease degradation is detected under these conditions, extended incubations and/or high concentrations of this enzyme may result in star activity. See the product FAQ for recommended reaction conditions for this enzyme.</p> <p>qPCR DNA Contamination (E. coli Genomic) A minimum of 10 units of WarmStart® Nt.BstNBI is screened for the presence of E. coli genomic DNA using SYBR® Green qPCR with primers specific for the E. coli 16S rRNA locus. Results are quantified using a standard curve generated from purified E. coli genomic DNA. The measured level of E. coli genomic DNA contamination is ≤ 1 E. coli genome.</p>	<p>Pass</p>

This product has been tested and shown to be in compliance with all specifications.

One or more products referenced in this document may be covered by a 3rd-party trademark. Please visit www.neb.com/trademarks for additional information.



Penghua Zhang
Production Scientist
16 Jun 2022



Michael Tonello
Packaging Quality Control Inspector
16 Jun 2022