

240 County Road Ipswich, MA 01938-2723 Tel 978-927-5054 Fax 978-921-1350 www.neb.com info@neb.com

## New England Biolabs Certificate of Analysis

Product Name: PaqCl®
Catalog Number: R0745S
Concentration: 10,000 U/ml

Unit Definition: One unit is defined as the amount of enzyme required to digest 1 µg

of Lambda DNA incubated for 1 hour at 37°C in a total reaction

volume of 50 μl.

Packaging Lot Number: 10144201
Expiration Date: 09/2023
Storage Temperature: -20°C

Storage Conditions: 300 mM NaCl, 10 mM Tris-HCl, 1 mM DTT, 0.1 mM EDTA, 50 %

Glycerol , 500 μg/ml BSA, (pH 7.4 @ 25°C)

Specification Version: PS-R0745S/L v1.0

PaqCI® Component List				
<b>NEB Part Number</b>	<b>Component Description</b>	Lot Number	Individual QC Result	
S0532SVIAL	PaqCI® Activator	10144206	Pass	
R0745SVIAL	PaqCI®	10144200	Pass	
B6004SVIAL	rCutSmart™ Buffer	10143286	Pass	

Assay Name/Specification	Lot # 10144201
Ligation and Recutting (Terminal Integrity) After a 10-fold over-digestion of Lambda DNA with PaqCl®, >95% of the DNA fragments can be ligated with T4 DNA ligase in 16 hours at 16°C. Of these ligated fragments, >95% can be recut with PaqCl®.	Pass
Functional Testing (15 minute Digest) A 50 μl reaction in CutSmart® Buffer containing 1 μg of Lambda DNA and 1 μl of PaqCl® incubated for 15 minutes at 37°C results in complete digestion as determined by agarose gel electrophoresis.	Pass
Exonuclease Activity (Radioactivity Release) A 50 ul reaction in CutSmart® Buffer containing 1 µg of a mixture of single and double-stranded [ ³H] E. coli DNA and a minimum of 100 units of PaqCl® incubated for 4 hours at 37°C releases <0.1% of the total radioactivity.	Pass
Endonuclease Activity (Nicking) A 50 µl reaction in CutSmart® Buffer containing 1 µg of supercoiled PhiX174 DNA and	Pass



R0745S / Lot: 10144201

Page 1 of 2

Assay Name/Specification	Lot # 10144201
a minimum of 30 units of PaqCl® incubated for 4 hours at 37°C results in <20% conversion to the nicked form as determined by agarose gel electrophoresis.	
Non-Specific DNase Activity (16 Hour) A 50 µl reaction in CutSmart® Buffer containing 1 µg of Lambda DNA and a minimum of 30 units of PaqCl® incubated for 16 hours at 37°C results in a DNA pattern free of detectable nuclease degradation as determined by agarose gel electrophoresis.	Pass
Protein Purity Assay (SDS-PAGE) PaqCl® is ≥ 95% pure as determined by SDS-PAGE analysis using Coomassie Blue detection.	Pass

This product has been tested and shown to be in compliance with all specifications.

One or more products referenced in this document may be covered by a 3rd-party trademark. Please visit www.neb.com/trademarks for additional information.

Penghaa Zhang Production Scientist

08 Apr 2022

Josh Hersey

Packaging Quality Control Inspector

08 Apr 2022



R0745S / Lot: 10144201

Page 2 of 2