NEW ENGLAND BioLabs Inc.
be INSPIRED drive DISCOVERY stay GENUINE
240 County Road
Ipswich, MA 01938-2723
Tel 978-927-5054
Fax 978-921-1350
www.neb.com
info@neb.com

New England Biolabs
Certificate of Analysis

Product Name:	*BamHI-HF®*
Catalog Number:	*R3136S*
Concentration:	*20,000 U/ml*
Unit Definition:	*One unit is defined as the amount of enzyme required to digest 1 µg of Lambda DNA in 1 hour at 37°C in a total reaction volume of 50 µl.*
Lot Number:	*10011443*
Expiration Date:	*01/2020*
Storage Temperature:	*-20°C*
Storage Conditions:	*50 mM KCl, 10 mM Tris-HCl (pH 7.4), 1 mM DTT, 0.1 mM EDTA, 50% Glycerol, 200 µg/ml BSA*
Specification Version:	*PS-R3136S/L v1.0*

BamHI-HF® Component List

NEB Part Number	Component Description	Lot Number	Individual QC Result
R3136SVIAL	BamHI-HF®	0101801	**Pass**
B7204SVIAL	CutSmart® Buffer	3081804	**Pass**
B7024SVIAL	Gel Loading Dye, Purple (6X)	0241804	**Pass**

Assay Name/Specification	Lot # 10011443
Exonuclease Activity (Radioactivity Release) A 50 µl reaction in CutSmart™ Buffer containing 1 µg of a mixture of single and double-stranded [^{3}H] E. coli DNA and a minimum of 100 units of BamHI-HF™ incubated for 4 hours at 37ºC releases <0.1% of the total radioactivity.	**Pass**
Endonuclease Activity (Nicking) A 50 µl reaction in CutSmart™ Buffer containing 1 µg of supercoiled PhiX174 DNA and a minimum of 100 Units of BamHI-HF™ incubated for 4 hours at 37ºC results in <10% conversion to the nicked form as determined by agarose gel electrophoresis.	**Pass**
Blue-White Screening (Terminal Integrity) A sample of pUC19 vector linearized with a 10-fold excess of BamHI-HF™, religated and transformed into an E. coli strain expressing the LacZ beta fragment gene results in <1% white colonies.	**Pass**
Ligation and Recutting (Terminal Integrity) After a 50-fold over-digestion of Lambda DNA with BamHI-HF™, >95% of the DNA fragments can be ligated with T4 DNA ligase in 16 hours at 16ºC. Of these ligated	**Pass**

NEW ENGLAND BioLabs Inc.
be INSPIRED drive DISCOVERY stay GENUINE
240 County Road
Ipswich, MA 01938-2723
Tel 978-927-5054
Fax 978-921-1350
www.neb.com
info@neb.com

Assay Name/Specification	Lot # 10011443
fragments, >95% can be recut with BamHI-HF™. **Non-Specific DNase Activity (16 Hour)** A 50 µl reaction in CutSmart™ Buffer containing 1 µg of Lambda DNA and a minimum of 100 Units of BamHI-HF™ incubated for 16 hours at 37ºC results in a DNA pattern free of detectable nuclease degradation as determined by agarose gel electrophoresis.	**Pass**

This product has been tested and shown to be in compliance with all specifications.

JianYing Luo
Production Scientist
07 Jun 2018

Michael Tonello
Packaging Quality Control Inspector
07 Jun 2018