

## New England Biolabs Certificate of Analysis

**Product Name:** *SacI-HF*<sup>®</sup>  
**Catalog Number:** R3156S  
**Concentration:** 20,000 U/ml  
**Unit Definition:** One unit is defined as the amount of enzyme required to digest 1 µg of Lambda DNA (*HindIII* digest) in 1 hour at 37°C in a total reaction volume of 50 µl.  
**Packaging Lot Number:** 10078404  
**Expiration Date:** 05/2022  
**Storage Temperature:** -20°C  
**Storage Conditions:** 100 mM NaCl, 10 mM Tris-HCl (pH 7.4), 1 mM DTT, 0.1 mM EDTA, 50% Glycerol, 200 µg/ml BSA  
**Specification Version:** PS-R3156S/L v2.0

SacI-HF <sup>®</sup> Component List			
NEB Part Number	Component Description	Lot Number	Individual QC Result
R3156SVIAL	SacI-HF <sup>®</sup>	10074526	Pass
B7204SVIAL	CutSmart <sup>®</sup> Buffer	10077115	Pass
B7024SVIAL	Gel Loading Dye, Purple (6X)	10075962	Pass

Assay Name/Specification	Lot # 10078404
<b>Endonuclease Activity (Nicking)</b> A 50 µl reaction in CutSmart <sup>™</sup> Buffer containing 1 µg of supercoiled PhiX174 DNA and a minimum of 60 Units of SacI-HF <sup>™</sup> incubated for 4 hours at 37°C results in <10% conversion to the nicked form as determined by agarose gel electrophoresis.	Pass
<b>Exonuclease Activity (Radioactivity Release)</b> A 50 µl reaction in CutSmart <sup>™</sup> Buffer containing 1 µg of a mixture of single and double-stranded [ <sup>3</sup> H] E. coli DNA and a minimum of 100 units of SacI-HF <sup>™</sup> incubated for 4 hours at 37°C releases <0.1% of the total radioactivity.	Pass
<b>Ligation and Recutting (Terminal Integrity)</b> After a 20-fold over-digestion of pXba DNA with SacI-HF <sup>™</sup> , >95% of the DNA fragments can be ligated with T4 DNA ligase in 16 hours at 16°C. Of these ligated fragments, >95% can be recut with SacI-HF <sup>™</sup> .	Pass
<b>Non-Specific DNase Activity (16 Hour)</b> A 50 µl reaction in CutSmart <sup>™</sup> Buffer containing 1 µg of Lambda-HindIII DNA and a	Pass

Assay Name/Specification	Lot # 10078404
minimum of 100 Units of SacI-HF™ incubated for 16 hours at 37°C results in a DNA pattern free of detectable nuclease degradation as determined by agarose gel electrophoresis.	
<b>Protein Purity Assay (SDS-PAGE)</b> SacI-HF™ is >95% pure as determined by SDS PAGE analysis using Coomassie Blue detection.	<b>Pass</b>
<b>Blue-White Screening (Terminal Integrity)</b> A sample of LITMUS28i vector linearized with a 10-fold excess of SacI-HF™, religated and transformed into an E. coli strain expressing the LacZ beta fragment gene results in <1% white colonies.	<b>Pass</b>

This product has been tested and shown to be in compliance with all specifications.

One or more products referenced in this document may be covered by a 3rd-party trademark. Please visit [www.neb.com/trademarks](http://www.neb.com/trademarks) for additional information.



\_\_\_\_\_  
 Penghua Zhang  
 Production Scientist  
 06 Aug 2020



\_\_\_\_\_  
 Michael Tonello  
 Packaging Quality Control Inspector  
 06 Aug 2020