

New England Biolabs Product Specification

<i>Product Name:</i>	<i>TaqI Methyltransferase</i>
<i>Catalog #:</i>	<i>M0219S</i>
<i>Concentration:</i>	<i>10,000 units/ml</i>
<i>Unit Definition:</i>	<i>One unit is defined as the amount of enzyme required to protect 1 µg Lambda DNA in 1 hour at 65°C in a total reaction volume of 20 µl against cleavage by TaqI restriction endonuclease.</i>
<i>Shelf Life:</i>	<i>24 months</i>
<i>Storage Temp:</i>	<i>-20°C</i>
<i>Storage Conditions:</i>	<i>100 mM NaCl, 10 mM Tris-HCl, 1 mM DTT, 0.1 mM EDTA, 50 % Glycerol, 200 µg/ml BSA, (pH 7.4 @ 25°C)</i>
<i>Specification Version:</i>	<i>PS-M0219S v1.0</i>
<i>Effective Date:</i>	<i>16 May 2018</i>

Assay Name/Specification (minimum release criteria)

Functional Testing (Methyltransferase) - A 20 µl reaction in CutSmart® Buffer supplemented with 80 µM SAM containing 1 µg of Lambda DNA and 1 unit of TaqI Methyltransferase incubated for 1 hour at 65°C followed by heat inactivation results in ≥ 95% protection from digestion with 10 units of TaqI in CutSmart® Buffer incubated at 65°C for 15 minutes as determined by agarose gel electrophoresis.

Exonuclease Activity (Radioactivity Release) - A 50 µl reaction in NEBuffer 2 containing 1 µg of a mixture of single and double-stranded [³H] *E. coli* DNA and a minimum of 100 units of TaqI Methyltransferase incubated for 4 hours at 65°C releases <0.1% of the total radioactivity.

Non-Specific DNase Activity (16 Hour) - A 50 µl reaction in NEBuffer 2 containing 1 µg of BstEII digested Lambda DNA and a minimum of 100 units of TaqI Methyltransferase incubated for 16 hours at 65°C results in a DNA pattern free of detectable nuclease degradation as determined by agarose gel electrophoresis.



Date 16 May 2018

Derek Robinson
Director of Quality Control

