

New England Biolabs Product Specification

<i>Product Name:</i>	<i>APE 1</i>
<i>Catalog #:</i>	<i>M0282S/L/V</i>
<i>Concentration:</i>	<i>10,000 units/ml</i>
<i>Unit Definition:</i>	<i>One unit is defined as the amount of enzyme required to cleave 20 pmol of a 34 mer oligonucleotide duplex containing a single AP site in a total reaction volume of 10 µl in 1 hour at 37°C.</i>
<i>Shelf Life:</i>	<i>30 months</i>
<i>Storage Temp:</i>	<i>-20°C</i>
<i>Storage Conditions:</i>	<i>10 mM Tris-HCl, 50 mM NaCl, 1 mM DTT, 0.05 mM EDTA, 200 µg/ml BSA, 50% Glycerol, (pH 8.0 @ 25°C)</i>
<i>Specification Version:</i>	<i>PS-M0282S/L v1.0</i>
<i>Effective Date:</i>	<i>19 Dec 2017</i>

Assay Name/Specification (minimum release criteria)

Endonuclease Activity (Nicking) - A 50 µl reaction in NEBuffer 4 containing 1 µg of supercoiled PhiX174 DNA and a minimum of 50 units of APE 1 incubated for 4 hours at 37°C results in <10% conversion to the nicked form as determined by agarose gel electrophoresis.

Exonuclease Activity (Radioactivity Release) - A 50 µl reaction in NEBuffer 4 containing 1 µg of a mixture of single and double-stranded [³H] *E. coli* DNA and a minimum of 50 units of APE 1 incubated for 4 hours at 37°C releases <0.5% of the total radioactivity.

Non-Specific DNase Activity (16 Hour) - A 50 µl reaction in NEBuffer 4 containing 1 µg of Lambda-HindIII DNA and a minimum of 50 units of APE 1 incubated for 16 hours at 37°C results in a DNA pattern free of detectable nuclease degradation as determined by agarose gel electrophoresis.

Protein Purity Assay (SDS-PAGE) - APE 1 is ≥ 95% pure as determined by SDS-PAGE analysis using Coomassie Blue detection.

One or more products referenced in this document may be covered by a 3rd-party trademark.
Please visit www.neb.com/trademarks for additional information.



Date 19 Dec 2017

Derek Robinson
Quality Approver

