

## New England Biolabs Product Specification

<i>Product Name:</i>	<i>T4 PDG (T4 Endonuclease V)</i>
<i>Catalog #:</i>	<i>M0308S/L</i>
<i>Concentration:</i>	<i>10,000 units/ml</i>
<i>Unit Definition:</i>	<i>One unit is defined as the amount of enzyme that catalyzes the conversion of 0.5 µg of UV-irradiated supercoiled pUC19 DNA to &gt;95% nicked plasmid in a total reaction volume of 20 µl in 30 minutes at 37°C. Nicking is assessed by agarose gel electrophoresis. Irradiated plasmid contains an average of 3-5 pyrimidine dimers.</i>
<i>Shelf Life:</i>	<i>24 months</i>
<i>Storage Temp:</i>	<i>-20°C</i>
<i>Storage Conditions:</i>	<i>10 mM Tris-HCl, 250 mM NaCl, 1 mM DTT, 0.1 mM EDTA, 50 % Glycerol, 0.15 % Triton® X-100, (pH 7.4 @ 25°C)</i>
<i>Specification Version:</i>	<i>PS-M0308S/L v1.0</i>
<i>Effective Date:</i>	<i>11 Jun 2018</i>

### Assay Name/Specification (minimum release criteria)

**Non-Specific DNase Activity (16 Hour)** - A 50 µl reaction in NEBuffer 2.1 containing 1 µg of Lambda-HindIII DNA and a minimum of 100 units of T4 PDG (T4 Endonuclease V) incubated for 16 hours at 37°C results in a DNA pattern free of detectable nuclease degradation as determined by agarose gel electrophoresis.

**Exonuclease Activity (Radioactivity Release)** - A 50 µl reaction in NEBuffer 2.1 containing 1 µg of a mixture of single and double-stranded [<sup>3</sup>H] *E. coli* DNA and a minimum of 30 units of T4 PDG (T4 Endonuclease V) incubated for 4 hours at 37°C releases <0.1% of the total radioactivity.

**Protein Purity Assay (SDS-PAGE)** - T4 PDG (T4 Endonuclease V) is ≥ 95% pure as determined by SDS-PAGE analysis using Coomassie Blue detection.



Date 11 Jun 2018

Derek Robinson  
Director of Quality Control

