

## New England Biolabs Product Specification

<i>Product Name:</i>	<i>Nt.AlwI</i>
<i>Catalog #:</i>	<i>R0627S/L</i>
<i>Concentration:</i>	<i>10,000 units/ml</i>
<i>Unit Definition:</i>	<i>One unit is defined as the amount of enzyme required to convert 1 µg of supercoiled pUC101 DNA (dam-/ dcm-) to open circular form in 1 hour at 37°C in a total reaction volume of 50 µl.</i>
<i>Shelf Life:</i>	<i>24 months</i>
<i>Storage Temp:</i>	<i>-20°C</i>
<i>Storage Conditions:</i>	<i>50 mM KCl, 10 mM Tris-HCl (pH 7.4), 1 mM DTT, 0.1 mM EDTA, 50% Glycerol, 200 µg/ml BSA</i>
<i>Specification Version:</i>	<i>PS-R0627S/L v2.0</i>
<i>Effective Date:</i>	<i>07 Mar 2014</i>

### Assay Name/Specification (minimum release criteria)

**Exonuclease Activity (Radioactivity Release)** - A 50 µl reaction in CutSmart™ Buffer containing 1 µg of a mixture of single and double-stranded [<sup>3</sup>H] *E. coli* DNA and a minimum of 50 units of Nt.AlwI incubated for 4 hours at 37°C releases <0.1% of the total radioactivity.

**Non-Specific DNase Activity (16 hour)** - A 50 µl reaction in CutSmart™ Buffer containing 1 µg of pUC101dam-dcm- DNA and a minimum of 10 units of Nt.AlwI incubated for 16 hours at 37°C results in a DNA pattern free of detectable nuclease degradation as determined by agarose gel electrophoresis. NOTE: although no nuclease degradation is detected under these conditions, extended incubations and/or high concentrations of this enzyme may result in star activity. See the product FAQ for recommended reaction conditions for this enzyme.

One or more products referenced in this document may be covered by a 3rd-party trademark.  
Please visit [www.neb.com/trademarks](http://www.neb.com/trademarks) for additional information.



Date 07 Mar 2014

Derek Robinson  
Quality Approver

